



Why are our toes so tiny? Walking, running and the evolution of a short forefoot in the genus Homo

Citation

Rolian, Campbell, Daniel E. Lieberman, John W. Scott. 2007. Why are our toes so tiny? Walking, running and the evolution of a short forefoot in the genus Homo. Abstracts of AAPA poster and podium presentations. American Journal of Physical Anthropology 132(S44): 202.

Published Version

<http://dx.doi.org/10.1002/ajpa.20577>

Permanent link

<http://nrs.harvard.edu/urn-3:HUL.InstRepos:2770519>

Terms of Use

This article was downloaded from Harvard University's DASH repository, and is made available under the terms and conditions applicable to Other Posted Material, as set forth at <http://nrs.harvard.edu/urn-3:HUL.InstRepos:dash.current.terms-of-use#LAA>

Share Your Story

The Harvard community has made this article openly available.
Please share how this access benefits you. [Submit a story](#).

[Accessibility](#)

Why are our toes so tiny? Walking, running and the evolution of a short forefoot in the genus *Homo*

Campbell Rolian, Daniel E. Lieberman, John W. Scott

Abstract:

Humans have an extremely short forefoot relative to total foot length. The derived pedal proportions of humans are thought to have evolved in the context of committed bipedalism, but the benefits of shorter toes for walking and/or running have not previously been tested. Short toes are typically associated with cursorial digitigrade mammals, where they improve the ability of the digital flexor apparatus – the muscles, tendons and ligaments that collectively flex and resist extension of the metatarsophalangeal (MTP) joints – to support the body and generate propulsion at the end of stance. We tested the hypothesis that in humans a shorter forefoot similarly improves locomotor performance by decreasing the force, power and work outputs of the digital flexor apparatus (DFA) during late stance, especially in running, when only one foot provides support and propulsion against high ground reaction forces. Kinematic, force and plantar pressure data were collected from a sample representing normal variation in toe length ($n=12$). Hindlimb kinematics, DFA force, power and work outputs were compared during barefoot walking and running in subjects with short, average and long forefeet in relation to body mass. Results suggest that individuals with relatively longer forefeet experience higher MTP joint moments, and their DFA generates more force, power and work than subjects with shorter forefeet, at both walking and running speeds. Contrary to our prediction, however, the difference between groups in DFA performance is not greater at running speeds. Implications for the evolution of endurance running in the genus *Homo* are discussed.